



# A NEW METHOD FOR THE MEASUREMENT OF THE BACKBITING RATE COEFFICIENT IN ACRYLATE RADICAL POLYMERIZATION

Yoshi W. Marien<sup>1</sup>, Paul H.M. Van Steenberge<sup>1</sup>, Katrin B. Kockler<sup>2,3</sup>, Christopher Barner-Kowollik<sup>2,3</sup>,

Marie-Françoise Reyniers<sup>1</sup>, Dagmar R. D'hooge<sup>1,4</sup>, Guy B. Marin<sup>1</sup>

<sup>1</sup> Laboratory for Chemical Technology  
Technologiepark 914, 9052 Zwijnaarde, Belgium

<http://www.lct.ugent.be> E-mail: dagmar.dhooge@ugent.be

<sup>3</sup> Institut für Biologische Grenzflächen  
Hermann-von-Helmholtz-Platz 1,  
76344 Eggenstein-Leopoldshafen, Germany

<sup>2</sup> Preparative Macromolecular Chemistry,  
Institut für Technische Chemie  
und Polymerchemie

Engesserstrasse 18, 76128 Karlsruhe, Germany

<sup>4</sup> Department of Textiles,  
Technologiepark 907, B-9052 Zwijnaarde, Belgium



## Introduction

Pulsed laser polymerization has emerged as the leading technique for the determination of the **propagation** rate coefficient  $k_p$ .<sup>[1]</sup> In the PLP-SEC trace, inflection points ( $L_j$ ;  $j = 1, 2, \dots$ ) are identified that are directly linked to  $k_p$  via:

$$L_j = k_p [M]_0 (j\Delta t)$$

For systems with several radical types, the obtained  $k_p$  must be seen as an apparent averaged one ( $k_{p,app}$ ). For example, in acrylate radical polymerization both end-chain radicals (ECRs;  $k_{p,ecr}$ ) and mid-chain radicals (MCRs;  $k_{p,mcr}$ ) can be present.<sup>[2,3]</sup> The tendency of ECRs to switch to MCRs is expressed by the **backbiting** rate coefficient  $k_{bb}$ .

In the present work, a **new and accurate method** is presented to determine  $k_{bb}$  from inflection point PLP-SEC data.


## Working principle of the new method

- Variation of the **solvent volume fraction** ( $\Phi_S=0-0.75$ )
  - Independent change of the average MCR lifetime
  - Increased sensitivity toward  $k_{bb}$
- Regression analysis using a detailed **kinetic Monte Carlo** model to inflection point data in the low frequency range


	Equation	Bulk ( $\Phi_S = 0$ )	Solution ( $\Phi_S = 0.75$ )
Dark time [s]	$\frac{1}{v}$	$10^{-2}$	$10^{-2}$
Average lifetime ECR [s]	$\frac{1}{k_{bb}}$	$10^{-3}$	$10^{-3}$
Average lifetime MCR [s]	$\frac{1}{k_{p,mcr}(1 - \Phi_S)[M]_{0,bulk}}$	$1.4 \cdot 10^{-3}$	$5.7 \cdot 10^{-3}$

## Kinetic Monte Carlo modeling for the simulation of the entire PLP-SEC trace



### Explicit calculation of the time dependence of the radical concentration



$E_{pulse} = 2.5$  mJ,  $\lambda = 351$  nm,  $[DMPA] = 5$  mmol L<sup>-1</sup>  
 $\Phi = 0.8$ ,  $\epsilon = 280$  L mol<sup>-1</sup> cm<sup>-1</sup>,  $V = 0.2$  mL,  $L = 5.2$  mm




### Differentiation in chain initiation capability




### Importance of short-long termination

Visualization of termination reactions



### Validation of the composite $k_t$ -model



#### Composite model

$$i \leq i_c: k_t^{app}(i, i) = k_t^{app}(1, 1) i^{-\alpha_S}$$
$$i > i_c: k_t^{app}(i, i) = k_t^{app}(1, 1) i_c^{-\alpha_S + \alpha_L} i^{-\alpha_L}$$


#### Simple power law model

$$k_t^{app}(i, i) = k_t^{app}(1, 1) i^{-\alpha}$$

**Experiment<sup>[4]</sup>:** PLP of *n*-butyl acrylate with DMPA as photoinitiator; frequency: 500 s<sup>-1</sup>


## In silico validation of the new method

**In silico** experiment:  $k_{p,app} = \hat{k}_{p,app}(k_{bb}^{input}) + e(\sigma)$  with  $e(\sigma)$  a standard normally distributed error



$\sigma = 300$  L mol<sup>-1</sup> s<sup>-1</sup>  
95% confidence interval:  $998 \pm 168$  s<sup>-1</sup>

→ **Very precise determination of  $k_{bb}$  in case of an acceptable experimental error**



$\sigma = 600$  L mol<sup>-1</sup> s<sup>-1</sup>  
95% confidence interval:  $959 \pm 280$  s<sup>-1</sup>

→ **Even for a large experimental error, an accurate value for  $k_{bb}$  results**

## Conclusions

- Experimental data** of the full PLP-SEC trace are accurately described via the **kMC technique**
- The different **chain initiation** capability of the DMPA fragments is confirmed and related to the peak intensities of the PLP-SEC trace
- Importance of **short-long termination** is demonstrated, indicating the possible extraction of short-long termination kinetics from PLP experiments
- PLP-SEC data can be used to test the validity of models for **apparent termination rate coefficients** at low monomer conversions
- Variation of the solvent volume fraction in the low frequency range allows to accurately determine  $k_{bb}$

## References

- [1] K.B. Kockler, A.P. Haehnel, T. Junkers, C. Barner-Kowollik, *Macromol. Rapid Commun.* **2016**, 37, 123.
- [2] B. Wenn, T. Junkers, *Macromol. Rapid Commun.* **2016**, 37, 781.
- [3] A.N. Nikitin, R.A. Hutchinson, M. Buback, P. Hesse, *Macromolecules* **2007**, 40, 8631.
- [4] C. Barner-Kowollik, F. Gunzler, T. Junkers, *Macromolecules* **2008**, 41, 8971.

## Acknowledgements

Y.W. M., P.H.M.V.S., M.-F.R., D.R.D., and G.B.M. acknowledge financial support from the Long Term Structural Methusalem Funding by the Flemish Government, the Interuniversity Attraction Poles Programme – Belgian State – Belgian Science Policy, and the Fund for Scientific Research Flanders (FWO; G.0065.13N). D.R.D. and P.H.M.V.S. acknowledge the FWO through a postdoctoral fellowship. C.B.-K. acknowledges continued support from the Karlsruhe Institute of Technology (KIT) in the context of the STN program of the Helmholtz association.